

# Treatment of Severe COVID-19 with Low-Dose Selinexor: Demonstration of Anti-Viral and Anti-Inflammatory Activities in a Randomized, International, Multicenter, Placebo-Controlled Phase 2 Clinical Trial

**George Geils Jr.**, Zainab Shahid, Don Stevens, Nathan C. Bahr, Jacek Skarbinski,  
Vaishali Sanchorawala, Marcelo Gareca, Ronda Oram, Miguel Marcos, Parameswaran Hari, Paul Boyce,  
Enrico Lallana, John P. Haran, Guenther Koehne, Otto Yang, Catherine B. Small, Ruben Niesvizky,  
Charles Brummitt, Piers Patten, Mansour Ceesay, Benjamin J. Gaborit, Ran Nir-Paz, Manish Sagar,  
Michael Kauffman, Sharon Shacham, Jatin Shah, Dayana Michel, Sharon Tamir, Tami Rashal, Lingling Li,  
Hong Yan, Yosef Landesman, Ralph A. Tripp, Thomas J. Walsh, Mezgebe Berhe



## Disclosures – George Geils Jr.


<b>Research Support</b>	None
<b>Consultant</b>	None
<b>Honoraria</b>	None
<b>Scientific Advisory Board</b>	None
<b>Major Stockholder</b>	None
<b>Speakers Bureau</b>	Amgen, Celgene, Janssen Oncology, and Sanofi Genzyme

# Exportin 1 (XPO1): Viral replication


- Exportin 1 (XPO1) mediates SARS-CoV-2 lifecycle and proinflammatory transcription factors
- XPO1 has been identified as a “hub” host protein for SARS-CoV propagation.
- XPO1 plays a central role in inflammation through regulation of the NF-kB and COX-2 pathways
- XPO1 facilitates the nuclear export of the HMGB1, RXR $\alpha$ , COMMD1, PPAR $\gamma$ , and GLTSCR2, all of which augment inflammatory signaling and inhibit innate immune response to enable viral infection

# Selinexor (XPOVIO®) XPO1 Inhibition: Unique Dual Mechanism of Action with Anti-Viral and Anti-inflammatory Activities


Class	Virus / Host Cellular Pathway	XPO1 Cargo	Outcome
<b>DIRECT - VIRAL XPO1 cargos essential for viral propagation</b>	SARS-CoV-2	ORF3b, ORF6, 9b, spike, N-protein	↓viral replication and propagation
	HIV	Rev	↓HIV late mRNA translation ↓virus production
	Influenza	NEP	↓viral replication
	HCMV	pp65/UL95	↓viral replication
	JCV	agnoprotein	↓viral RNA processing
<b>INDIRECT – HOST XPO1 cargos. Nuclear entrapment reduces inflammation &amp; blocks viral propagation</b>	Cell membrane receptor	ACE-2	↓SARS-CoV-2 viral propagation
	NF-κB	IκB, COMMD1, FoxO	↓NFκB gene expression ↑oxidative stress response
	IL1β	RXRα	↓inflammatory response
	COX2, iNOS	COX2 and iNOS mRNAs	↓inflammatory mediators
	PPARγ	PPARγ	↑anti-inflammatory and cytoprotective response
	TLR2, TLR4, RAGE	HMGB1	↓necrosis induced inflammation
	FoxO, FoxP	Forkhead Proteins	↓inflammatory response
	HIF-1	COMMD1	↓NFκB gene expression
	Nrf2	Nrf2	↑oxidative stress response
	Rig1 Innate Immunity	GLTSCR2	↑RIG-I activity ↑interferon β production
	IRF7	KSHV ORF 45	↑IFNα/β production
	p53	HPV E2 and E6	↓p53 degradation ↓apoptotic response



# Selinexor Inhibits SARS-CoV-2 Viral Propagation and Shedding and Induces Nuclear Accumulation of ACE2 *In Vitro*



# Selinexor Decreases SARS-CoV-2 Viral Load and Severity of Rhinitis and Lung Inflammation *In Vivo*



# XPORT-CoV-1001: A Phase 2 Randomized Single-Blind Study to Evaluate the Activity and Safety of Low Dose Oral Selinexor (KPT-330) in Patients with Severe COVID-19

*~70 International Study Sites, ~8 countries*  
*2 interim analyses (~74 randomized and ~124 randomized)*

2:1 → 1:1  
randomization  
**N ~ 223**

Hospitalized  
Patients ≥18  
years old with  
COVID-19

## Oral Selinexor

20 mg Days 1, 3, and 5 of each week  
for up to 2 weeks

If the patient is tolerating therapy well and clinically  
benefitting, dosing can continue for additional 2  
weeks on Days 15, 17, 19, 22, 24, 26

## Oral Placebo

Days 1, 3, and 5 of each week  
for up to 2 weeks

## Primary endpoints:

### Day 14 Ordinal Scale Improvement (OSI) -

Proportion of patients with at least a 2-point improvement  
in the Ordinal Scale from baseline to Day 14

### Key Secondary endpoint

- Overall death rate on Day 28
- Rate of mechanical ventilation
- Time to mechanical ventilation
- Time to an improvement of 2 points using Ordinal Scale Improvement (TTCI-2)



# XPORT-CoV-1001: Inclusion Criteria

Eligible patients had symptoms of severe COVID-19 as demonstrated by:

## **At least 1 of the following:**

- Fever
- Cough
- Sore throat
- Malaise
- Headache
- Muscle pain
- Shortness of breath at rest or with exertion
- Confusion
- Symptoms of severe lower respiratory symptoms including dyspnea at rest or respiratory distress

**AND**

## **Clinical signs indicative of lower respiratory infection with COVID-19, with at least 1 of the following:**

- SpO2  $\leq$ 92% or requires  $\geq$ 4 LPM oxygen by nasal canula, or
- Non-rebreather/Ventimask (or similar device) or
- High-flow nasal canula in order to maintain SpO2  $\geq$ 92%.

Patients with COPD or chronic lung disease must demonstrate evidence of increased oxygen needs above baseline

# XPORT-CoV-1001: Patient Characteristics (ITT)

CHARACTERISTIC	N = 117
<b>Age, years</b>	
Median, (range)	55.0 (25-98)
<b>Sex, n (%)</b>	
Male	68 (58.1)
<b>Concomitant Therapies, n (%)</b>	
Remdesivir	59 (50.4)
Hydroxychloroquine	1 (0.9)
Systemic corticosteroid	54 (46.2)
IL-6 inhibitor*	10 (8.6)
<b>No. of High-Risk Comorbidities, n (%)</b>	
≥2	43 (36.8)
1	26 (22.2)
None	46 (39.3)
<b>Comorbidities, n (%)</b>	
Hypertension	44 (37.6)
Metabolic	87 (74.4)
Cancer	13 (11.1)
Diabetes	44 (37.6)
Cardiac	26 (22.2)
Renal	22 (18.8)




# XPORT-CoV-1001: Treatment-Related Adverse Events (ITT)

n, (%)	Selinexor N=65				Placebo N=50			
	Grade 1/2	Grade 3	Grade 4	Total	Grade 1/2	Grade 3	Grade 4	Total
<b>Any Treatment-Related Adverse Event</b>	14 (21.5)	4 (6.2)	1 (1.5)	19 (29.2)	8 (16.0)	2 (4.0)	0 (0)	10 (20.0)
Hematological								
<b>Hyponatremia</b>	<b>3 (4.6)</b>	<b>1 (1.5)</b>	<b>0 (0)</b>	<b>4 (6.2)</b>	<b>3 (6.0)</b>	<b>0 (0)</b>	<b>0 (0)</b>	<b>3 (6.0)</b>
Neutropenia	1 (1.5)	0 (0)	0 (0)	1 (1.5)	0 (0)	0 (0)	1 (2.0)	1 (2.0)
<b>Non-Hematological</b>								
Nausea	5 (7.7)	0 (0)	0 (0)	5 (7.7)	4 (8.0)	0 (0)	0 (0)	4 (8.0)
Constipation	1 (1.5)	0 (0)	0 (0)	1 (1.5)	2 (4.0)	0 (0)	0 (0)	2 (4.0)
Cough	2 (3.1)	0 (0)	0 (0)	2 (3.1)	0 (0)	0 (0)	0 (0)	0 (0)
Headache	0 (0)	0 (0)	0 (0)	0 (0)	2 (4.0)	0 (0)	0 (0)	2 (4.0)
Vomiting	1 (1.5)	0 (0)	0 (0)	1 (1.5)	1 (2.0)	0 (0)	0 (0)	1 (2.0)
<b>Serious TRAEs</b>				<b>2 (3.1)</b>				<b>0 (0)</b>

# XPORT-CoV-1001: Efficacy (ITT)

Parameter	Selinexor N=66 n (%)	Placebo N=51 n (%)
Clinical improvement Day 14 (2-point improvement in ordinal scale), n (%)	35 (53.0)	28 (54.9)
Time to clinical improvement (2 points), median days (range)	11.0 (9.0, 16.0)	9.0 (8.0, 16.0)
Overall Mortality at Day 28	10 (15.2)	2 (3.9)
Mechanical ventilation required, n (%)	9 (13.6)	5 (9.8)
Admission to ICU, n (%)	33 (50.0)	23 (45.1)
Length of hospitalization, days, median (range)	9.0 (2-39)	8.0 (3-42)
Hospital discharge, n (%)	48 (72.7)	39 (76.5)
Conversion to negative SARS-CoV-2 PCR, n (%)	24 (36.4)	10 (19.6)



## XPORT-CoV-1001: Safety Outcomes (Low LDH/DD)

Parameter	Low LDH/DD*		High LDH/DD*	
	Selinexor N=38 n (%)	Placebo N=27 n (%)	Selinexor N=27 n (%)	Placebo N=23 n (%)
Patients with at least 1 AE	24 (63.2)	14 (51.9)	21 (77.8)	12 (52.2)
Patients with at least 1 Grade 3/4 AE	3 (7.9)	1 (3.7)	6 (22.2)	7 (30.4)
SAEs	5 (13.2)	1 (3.7)	11 (40.7)	5 (21.7)
Non-fatal SAEs	3 (7.9)	0 (0)	2 (7.4)	3 (13)
Invasive MV or ECMO	3 (7.9)	3 (11.1)	6 (22.2)	3 (13.0)
Death within 30 days of last dose	2 (5.3)	1 (3.7)	9 (33.3)	2 (8.7)

\* The Low LDH/DD subgroup was comprised of patients with LDH ≤370 U/L or D-dimer ≤600 mcg/L FEU. The High LDH/DD subgroup was comprised of patients with LDH >370 U/L and D-dimer >600 mcg/L FEU.

- No related Grade 5 AEs were reported in the mITT population
- Side effects were generally reversible and managed with dose modifications and/or standard supportive care

# XPORT-CoV-1001: Efficacy – Selinexor improved clinical status as compared with placebo (Low LDH/DD)

Parameter	Low LDH/DD*		High LDH/DD*	
	Selinexor N=38 n (%)	Placebo N=28 n (%)	Selinexor N=28 n (%)	Placebo N=23 n (%)
Hospital Discharged by Day 14	30 (78.9)	16 (57.1)	13 (46.4)	14 (60.9)
OSI-2 by Day 14	30 (78.9)	18 (64.3)	12 (42.9)	15 (65.2)






\* The Low LDH/DD subgroup was comprised of patients with LDH  $\leq 370$  U/L or D-dimer  $\leq 600$  mcg/L FEU. The High LDH/DD subgroup was comprised of patients with LDH  $> 370$  U/L and D-dimer  $> 600$  mcg/L FEU.

# XPORT-CoV-1001: Efficacy – Selinexor improved PCR negative conversion rate and time to PCR negative as compared with placebo (Low LDH/DD)

Parameter	Low LDH/DD*		High LDH/DD*	
	Selinexor N=38 n (%)	Placebo N=28 n (%)	Selinexor N=28 n (%)	Placebo N=23 n (%)
Converted to PCR negative status	16 (42.1)	8 (28.6)	11 (39.3)	7 (30.4)

\* The Low LDH/DD subgroup was comprised of patients with LDH  $\leq 370$  U/L or D-dimer  $\leq 600$  mcg/L FEU. The High LDH/DD subgroup was comprised of patients with LDH  $> 370$  U/L and D-dimer  $> 600$  mcg/L FEU.

# XPORT-CoV-1001: Selinexor reduces cytokines associated with COVID-19 (Low LDH/DD)



**ARM**

— Selinexor  
— Placebo

Parameter	Selinexor N=38 n (%)	Placebo N=28 n (%)
LDH	11/34 (32)	3/24 (13)
D-dimer	21/34 (62)	13/25 (52)
CRP	19/38 (50)	12/25 (48)

# XPORT-CoV-1001: Summary and Conclusions

- XPO1 is an important target with and critical role in viral replication cycle
- Selinexor has demonstrated anti-viral activity in vitro and in vivo
  - Selinexor protects healthy cells from infection via reduction in ACE receptor expression on cell surface
- Oral selinexor has a dual role in COVID-19, with both an anti-viral and anti-inflammatory
  - ITT: Improvement in converted to PCR negative status: from 19.6% to 36.4%
- Clinical Benefit in subset of patients with normal LDH or  $< 2$ ULN of D Dimer
  - Improvement in Hospital Discharge by Day 14: 57.1% to 78.9%
  - Improvement in OSI-2 by Day 14: 64.3% to 78.9%
  - Converted to PCR negative status: 28.6% to 42.1%
- Future studies to evaluate the safety and efficacy of oral selinexor in hospitalized patients with confirmed severe COVID-19 and Low LDH/DD as well as in patients with mild/moderate COVID-19 and Outpatient COVID are warranted



# Acknowledgements

## Patients, their Families, and Caregivers

## Investigators, Co-investigators and Study teams at each participating center

- Roper Hospital, Charleston, SC
- Levine Cancer Institute, Atrium Health, Charlotte NC
- Norton Healthcare, Louisville KY
- University of Kansas Medical Center, Kansas City, KS
- Kaiser Permanente Oakland, Oakland, CA
- Boston Medical Center, Boston, MA
- Lehigh Valley Hospital, Allentown PA
- Advocate Christ Medical Center Oak Lawn, IL
- University Hospital of Salamanca-University of Salamanca-IBSAL, Salamanca, Spain
- Medical College of Wisconsin, Milwaukee, WI
- Northside Hospital, Atlanta GA
- Kaiser Permanente Sacramento, Sacramento, CA
- University of Massachusetts Medical School, Worcester, MA
- Miami Cancer Institute at Baptist Health, Miami FL
- University of California Los Angeles, Los Angeles, CA
- Weill Cornell Medicine, New York, NY
- Aurora St. Luke's Medical Center, Milwaukee, WI
- Kings College Hospital, London UK
- Princess Royal University Hospital, Kent, UK
- Hotel-Dieu Hospital and INSERM CIC 1413, Nantes University Hospital, Nantes, France
- Hadassah Medical Center, Jerusalem, Israel
- Karyopharm Therapeutics, Newton MA
- University of Georgia College of Veterinary Medicine, Athens, GA
- Baylor Scott & White Dallas, Dallas TX

***This study was sponsored by Karyopharm Therapeutics***

Thank You

